



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

November 7, 2016

Michael D. Hall
Product Registration Manager
BASF Corporation, Agricultural Products
P.O. Box 13528, 26 Davis Drive
Research Triangle Park, NC 27709

Subject: Label Amendment – revised master label and proposed supplemental label
Product Name: Coronet Fungicide Seed Treatment
EPA Registration Number: 7969-274
Application Date: 7/21/2016
Decision Number: 519664

Dear Michael D. Hall:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. "To distribute or sell" is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company's website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product's label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA's Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 7969-274
Decision No. 519664

with FIFRA section 6. If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at mellor.nathan@epa.gov.



Shaja B. Joyner, Product Manager 20
Fungicide-Herbicide Branch
Registration Division 7505P



We create chemistry

Group 7 | 11 Fungicide

ACCEPTED
11/07/2016
Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 7969-274

Coronet®

Fungicide Seed Treatment

For use in arugula, brassicas, bulb vegetables, cotton, cucurbits, legume vegetables, rapeseed, soybean and sunflower

Active Ingredients*:

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl]methoxy-, methyl ester) 9.0%

boscalid: 3-pyridinecarboxamide, 2-chloro-N-(4-chloro(1,1'-biphenyl)-2-yl)- 18.0%

Other Ingredients: 73.0%

Total: 100.0%

* Contains 0.83 pound of pyraclostrobin and 1.67 pounds of boscalid per gallon.

EPA Reg. No. 7969-274

EPA Est. No.

KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

See inside for complete First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty, and state-specific crop and/or use site restrictions.

In case of an emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

FIRST AID

If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything to an unconscious person.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after first 5 minutes; then continue rinsing eyes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION. Harmful if swallowed. Avoid contact with skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long-sleeved shirt and long pants, socks, shoes, and chemical-resistant gloves. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.

Personal Protective Equipment (PPE)

A dust-mist filtering respirator with MSHA/NIOSH approval number prefix TC-21C or a NIOSH-approved respirator with any N, R, P, or HE filter is required when using this product.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material (such as nitrile, butyl, neoprene and/or barrier laminate)
- Shoes plus socks

Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exists, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

Environmental Hazards

This pesticide is toxic to fish and aquatic invertebrates. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

DO NOT discharge effluent containing this product into lakes, streams, ponds, estuaries, oceans, or other waters unless in accordance with the requirements of a National Pollutant Discharge Elimination System (NPDES) permit and the permitting authority has been notified in writing prior to discharge. **DO NOT** discharge effluent containing this product to sewer systems without previously notifying the local sewage treatment plant authority. For guidance, contact your State Water Board or Regional Office of the EPA.

For Terrestrial Uses. DO NOT apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. **DO NOT** contaminate water when disposing of equipment washwater or rinsate.

Surface Water Advisory

This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water.

This product is classified as having medium potential for reaching both surface water and aquatic sediment via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of boscalid from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall or irrigation is forecasted to occur within 48 hours.

Ground Water Advisory

This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

Directions For Use

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

DO NOT apply this product in a way that will contact workers or other persons. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

STORAGE AND DISPOSAL

DO NOT contaminate water, food, or feed by storage or disposal.

Pesticide Storage

Store in original containers only. Keep container closed when not in use. **DO NOT** store near food or feed. In case of spill on floor or paved surfaces, mop and remove to chemical waste storage area until proper disposal can be made if product cannot be used according to label.

Pesticide Disposal

Wastes resulting from using this product may be disposed of on-site or at an approved waste disposal facility. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container Handling

Nonrefillable Container. DO NOT reuse or refill this container. Triple rinse or pressure rinse container (or equivalent) promptly after emptying; then offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

Triple rinse containers small enough to shake (capacity ≤ 5 gallons) as follows: Empty the remaining contents into application equipment or a mix tank and drain for 10 seconds after the flow begins to drip. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank, or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times.

Pressure rinse as follows: Empty the remaining contents into application equipment or mix tank and continue to drain for 10 seconds after the flow begins to drip. Hold container upside down over application equipment or mix tank, or collect rinsate for later use or disposal. Insert pressure rinsing nozzle in the side of the container and rinse at about 40 PSI for at least 30 seconds. Drain for 10 seconds after the flow begins to drip.

In Case of Emergency

In case of large-scale spillage regarding this product, call:

- CHEMTREC 1-800-424-9300
- BASF Corporation 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation 1-800-832-HELP (4357)

Steps to be taken in case this material is released or spilled:

- Dike and contain spill with inert material (sand, earth, etc.) and transfer liquid and solid diking material to separate containers for disposal.
- Remove contaminated clothing and wash affected skin areas with water.
- Wash clothing before reuse.
- Keep spill out of all sewers and open bodies of water.

Product Information

Coronet® fungicide seed treatment, a suspension concentrate (SC), is a broad-spectrum fungicide that provides preventive seed and seedling protection against seedborne fungi causing seed decay and the soilborne pathogens *Rhizoctonia solani* and *Penicillium* spp. **Coronet** may be applied in conjunction with applications of other registered seed treatment and crop protection products. Make all applications according to the use directions in this label. Use the higher rates of **Coronet** when disease pressure is expected to be high.

Resistance Management

The active ingredients in **Coronet**, boscalid and pyraclostrobin, belong to the groups of respiration inhibitors classified by the EPA as **Carboxamides** and **Quinone Outside Inhibitors (QoI)** or **Target Site of Action Group 7** and **Target Site of Action Group 11** fungicides, respectively. Due to the minimal amounts of active ingredient in **Coronet** applied to the seed that relocate to the upper parts of the plant, the potential for development of resistance to both fungicides is extremely low. Seed treatments of **Coronet** can, therefore, be followed by foliar use and resistance management guidelines of foliar use fungicides containing boscalid, pyraclostrobin, or both (**Cabrio® EG fungicide**, **Endura® fungicide**, **Headline® fungicide** or **Pristine® fungicide**) as specified on the respective labels.

Application Instructions

For use on “crop” seed at commercial treatment facilities.

Not for use on farm.

Apply **Coronet** as a water-based slurry using standard slurry or mist-type seed treatment application equipment. The exact amount of water needed to provide the slurry rate (ml/100 kg or fl ozs/cwt of seed) for optimum coverage is difficult to predict because weather conditions (hot or cold), seed type and surface, and equipment used all have a bearing on coverage. Consult a seed treatment specialist regarding slurry rates required for the crop seed to be treated with **Coronet**. Mix the required amount of **Coronet** with sufficient water to provide uniform and complete coverage on the seed surface. Uniform application on seed and complete seed coverage are necessary for seed safety and best disease protection.

Seed should be of good quality and well cleaned prior to treatment. The purchaser of **Coronet** is responsible for ensuring that all seed treated with this product are

adequately dyed with a suitable color to prevent its accidental use as food for man or feed for animals. Refer to 21CFR, Part 2.25. Use an EPA-approved dye or colorant that imparts an unnatural color to the seed. Any dye or colorant added to treated seed must be cleared for use under 40CFR, Part 180.900, “**Exemptions from the requirement of a tolerance**”. Federal regulations have established official tolerances or exemptions from tolerances for residues on food and forage crops that should not be exceeded when the product (dye or colorant) is used at required rates.

Consult the seed treatment specialist for calibration and operation procedures of the seed treatment equipment being used. Mechanical agitation is required for proper mixing of **Coronet**. Prepare no more mixture of **Coronet** than is needed for immediate treating. Add 3/4 of the required volume of water to the mix tank. With agitation running, add the required amount of **Coronet** in the tank along with other seed treatment components.

Add the remaining 1/4 of the required water. Follow the mixing sequence listed below. Agitate thoroughly before and during application.

Coronet With Tank Mix Partners

Coronet mixes easily with water and other water-based seed treatments manufactured by BASF Corporation. When using **Coronet** in a tank mixture with other seed treatment products, observe all directions for use, crops/sites, use rates, dilution ratios, precautions, and limitations that appear on the tank mix partner label. **DO NOT** exceed label dosage or application rate, and follow the most restrictive label precautions and limitations. **DO NOT** mix with any product that prohibits such mixing. Make all applications according to the use directions.

When mixing **Coronet** with products from other manufacturers, test compatibility prior to use by conducting a jar test as follows:

1. Mix all intended seed treatments with the appropriate amount of water in a clear glass container.
2. Mix well and allow to sit for 1 hour.
3. Remix and observe for incompatibility.

Physical compatibility is only part of compatibility. Testing also needs to be done with the seed to make sure the combination of products does not have detrimental effects on the seed.

Tank Mixing Instructions

1. Add 3/4 of the required amount of water to the mix tank and turn on the agitation. Mechanical agitation is preferred.
2. Follow the following mixing sequence: dye or colorant, wettable powders, soluble powders, flowable formulations, emulsifiable concentrates, and remaining 1/4 of the required water. The 1/4 of the required water as the last item for addition to the premix tank can be used as rinsate for the chemical container or to adjust the total slurry to compensate for water displacement by powdered products.

3. Maintain agitation until the entire slurry mixture has been used.
4. Ensure that seed is dry prior to bagging.

Seed treated with this product that is then packaged or bagged for future use must contain the following labeling on the outside of the seed package or bag.

Seed Labeling Note

“This seed has been treated with **Coronet® fungicide seed treatment** containing boscalid and pyraclostrobin. **DO NOT** use treated seed for food, feed or oil purposes. Store treated seed away from food and feedstuffs. **DO NOT** allow children, pets or livestock to have access to treated seed. Wear long pants, longsleeved shirt, shoes and socks, and chemical resistant gloves made of any waterproof material when opening this bag or handling (e.g., loading, pouring) treated seed or seed pieces. Treated seed exposed on soil surfaces may be hazardous to wildlife. Plant treated seed into the soil to the recommended minimum depth or greater to minimize exposure. Cover or collect treated seed spilled during loading and planting, in particular at row ends and field corners. Dispose of all excess treated seed by burying seed away from bodies of water. **DO NOT** enter or allow worker entry into areas with treated seeds during the restricted entry interval of 12 hours, with the exception that workers may enter if they will have no contact with the soil/media sub-surface. **DO NOT** contaminate bodies of water when disposing of planting equipment washwater. Dispose of seed packaging or containers in accordance with local requirements. Excess treated seed may be used for ethanol production only if (1) by-products are not used for livestock feed and (2) no measurable residues of pesticide remain in ethanol by-products that are used in agronomic practice. In the event of crop failure or harvest of a crop grown from this seed, the field may be replanted immediately with alfalfa, berries, brassica vegetables, bulb vegetables, celery, cotton, cucurbits, dry beans, fruiting vegetables, grape, leafy vegetables, leafy petioles, legume vegetables (except cowpea, field pea, and grain lupin), low growing berry, oilseed crops, peanut, pome fruits, rapeseed (includes canola and crambe), soybean, spinach, stone fruits, strawberry, sunflower, tree nuts, turnip greens, or any other crop for which tolerances for both boscalid and pyraclostrobin exist. For all other crops, the minimum plant back interval is 12 months from the date this seed was planted. A cover crop without tolerances for both boscalid and pyraclostrobin may be planted sooner than the 12 month interval for purposes of erosion control or soil improvement; however, the crop may not be grazed or harvested for food or feed.”

Coronet® fungicide seed treatment Crop-specific Use Directions

Crop	Target Disease	Product Use Rate ml/100 kg seed (gram ai/100 kg seed)	Product Use Rate fl ozs/100 lbs seed (lb ai/100 lbs seed)
Arugula	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> , <i>Phoma lingam</i> and other seedborne fungi causing seed decay and seedling damping off	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)
Brassica Vegetables	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> , <i>Phoma lingam</i> and other seedborne fungi causing seed decay and seedling damping off	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)
Bulb Vegetables	Control of: Seed and seedling disease caused by seedborne fungi (<i>Botrytis</i> , <i>Aspergillus</i> and <i>Penicillium</i> spp.) causing seed decay and seedling damping off	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)
Cotton¹	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)
Cucurbits¹	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)
Legume vegetables (dried shelled peas and beans, except cowpea, field pea, and grain lupin) Broad bean, Chickpea, Fava bean, Guar, Lablab bean, Lentil, Pigeon pea <i>Lupinus</i> spp.: Sweet lupin, White lupin, White sweet lupin	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off	52.2 to 202 (15.6 to 60.6)	0.8 to 3.1 (0.015 to 0.06)

(continued)

Coronet® fungicide seed treatment Crop-specific Use Directions (continued)

Crop	Target Disease	Product Use Rate ml/100 kg seed (gram ai/100 kg seed)	Product Use Rate fl ozs/100 lbs seed (lb ai/100 lbs seed)
<p>Legume vegetables (dried shelled peas and beans, except cowpea, field pea, and grain lupin) (continued)</p> <p><i>Phaseolus</i> spp.: Black bean, Field bean, Kidney bean, Lima bean, Navy bean, Pink bean, Pinto bean, Tepary bean</p> <p><i>Vigna</i> spp.: Adzuki bean, Black-eyed pea, Catjang, Crowder pea, Moth bean, Mung bean, Rice bean, Southern pea, Urd bean</p>	<p>Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off</p>	<p>52.2 to 202 (15.6 to 60.6)</p>	<p>0.8 to 3.1 (0.015 to 0.06)</p>
<p>Rapeseed (cultivars, varieties and/or hybrids, including canola and crambe)</p>	<p>Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i>, <i>Phoma lingam</i> and other seedborne fungi causing seed decay and seedling damping off</p>	<p>202 to 404 (60.6 to 121.1)</p>	<p>3.1 to 6.2 (0.06 to 0.12)</p>
<p>Sobean (<i>Glycine</i> spp.)</p>	<p>Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off</p>	<p>52.2 to 202 (15.6 to 60.6)</p>	<p>0.8 to 3.1 (0.015 to 0.06)</p>
<p>Sunflower</p>	<p>Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> and other seedborne fungi causing seed decay and seedling damping off</p>	<p>104 to 202 (31.2 to 60.6)</p>	<p>1.6 to 3.1 (0.03 to 0.06)</p>
<p>¹ For control of seed and seedling disease (damping off) caused by <i>Pythium</i> spp., use Coronet only in conjunction with labeled rates of mefenoxam- or metalaxyl-containing seed treatment products.</p>			

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as the presence of other or untested or unapproved materials, seed of low quality or low vigor or low germination, use of the product in a manner inconsistent with its labeling, misapplication of the product, or weather conditions at planting or environmental conditions during seed storage, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

ST1108

***Cabrio, Coronet, Endura, Headline and Pristine** are registered trademarks of BASF.*

© 2016 BASF Corporation
All rights reserved.

007969-00274.20160601d.**NVA 2016-04-314-0119**

Supersedes: NVA 2009-04-314-0155

Supplemental: NVA 2016-04-314-0132

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry

Supplemental Label

Coronet®

Fungicide Seed Treatment

For use on arugula

This supplemental label expires on November 7, 2019, and must not be used or distributed after this date. **Active Ingredients*:**

pyraclostrobin: (carbamic acid, [2-[[[1-(4-chlorophenyl)-1H-pyrazol-3-yl]oxy]methyl]phenyl] methoxy-, methyl ester)	9.0%
boscalid: 3-pyridinecarboxamide, 2-chloro-N-(4-chloro(1,1'-biphenyl)-2-yl)-	18.0%
Other Ingredients:	<u>73.0%</u>
Total:	100.0%

* Contains 0.83 pound of pyraclostrobin and 1.67 pounds of boscalid per gallon.

EPA Reg. No. 7969-274


Directions For Use

- It is a violation of federal law to use this product in a manner inconsistent with its labeling.
- The supplemental labeling and the entire **Coronet® fungicide seed treatment** container label, EPA Reg. No. 7969-274, must be in possession of the user at the time of application.
- Read the label affixed to the container for **Coronet** before applying.
- Use of **Coronet** according to this labeling is subject to the use precautions and limitations imposed by the label affixed to the container for **Coronet**.

Application Instructions

Refer to the **Coronet** container label for specific seed treatment instructions and seed container labeling requirements.

For use on “crop” seed at commercial treatment facilities. **Not for use on Farm.**



Coronet® fungicide seed treatment Crop-specific Use Directions

Crop	Target Disease	Product Use Rate ml/100 kg seed (gram ai/100 kg seed)	Product Use Rate fl ozs/100 lbs seed (lb ai/100 lbs seed)
Arugula	Control of: Seed and seedling disease caused by <i>Rhizoctonia solani</i> , <i>Phoma lingam</i> and other seedborne fungi causing seed decay and seedling damping off.	202 to 404 (60.6 to 121.1)	3.1 to 6.2 (0.06 to 0.12)

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as the presence of other or untested or unapproved materials, seed of low quality or low vigor or low germination, use of the product in a manner inconsistent with its labeling, misapplication of the product, or weather conditions at planting or environmental conditions during seed storage, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BUYER'S EXCLUSIVE REMEDY AND BASF'S EXCLUSIVE LIABILITY, WHETHER IN CONTRACT, TORT, NEGLIGENCE, STRICT LIABILITY, OR OTHERWISE, SHALL BE LIMITED TO REPAYMENT OF THE PURCHASE PRICE OF THE PRODUCT.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BASF AND THE SELLER DISCLAIM ANY LIABILITY FOR CONSEQUENTIAL, EXEMPLARY, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF. ST1108

Coronet is a registered trademark of BASF.

© 2016 BASF Corporation
All rights reserved.

007969-00274.20160601b.NVA 2016-04-314-0132

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709



We create chemistry